

Please check whether you have the right question paper.

- N.B.: 1) Questions No. 1 is compulsory.
2) Answer any three from remaining five questions.

1. a) If $\tan \frac{x}{2} = \tan h \frac{u}{2}$, show that $u = \log \left[\tan \left(\frac{\pi}{4} + \frac{x}{2} \right) \right]$. [3]

b) Prove that the following matrix is orthogonal & hence find A^{-1} , [3]

$$A = \frac{1}{3} \begin{bmatrix} -2 & 1 & 2 \\ 2 & 2 & 1 \\ 1 & -2 & 2 \end{bmatrix}$$

c) State Euler's theorem on Homogeneous function of two variables & if [3]

$$u = \frac{x+y}{x^2+y^2} \text{ then evaluate } x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$$

d) If $u = r^2 \cos 2\theta$, $v = r^2 \sin 2\theta$. Find $\frac{\partial(u, v)}{\partial(r, \theta)}$. [3]

e) Find the n^{th} derivative of $\cos 5x \cdot \cos 3x \cdot \cos x$. [4]

f) Evaluate: $\lim_{x \rightarrow 0} \left(\frac{2x+1}{x+1} \right)^{\frac{1}{x}}$. [4]

2. a) Solve $x^4 - x^3 + x^2 - x + 1 = 0$. [6]

b) If $y = e^{\tan^{-1} x}$. Prove that [6]

$$(1+x^2)y_{n+2} + [2(n+1)x-1]y_{n+1} + n(n+1)y_n = 0.$$

c) Examine the function $f(x, y) = xy(3-x-y)$ for extremes values & [8]
also find maximum and minimum values of $f(x, y)$.

TURN OVER

3. a) Investigate for what values of λ & μ the equations $x+y+z=6$; [6]

$$x+2y+3z=10; x+2y+\lambda z=\mu \text{ have}$$

- i) no solution,
- ii) a unique solution,
- iii) infinite no. of solutions.

b) If $u = f\left(\frac{y-x}{xy}, \frac{z-x}{xz}\right)$, show that $x^2 \frac{\partial u}{\partial y} + y^2 \frac{\partial u}{\partial z} + z^2 \frac{\partial u}{\partial x} = 0$. [6]

c) Prove that $\log\left(\frac{a+ib}{a-ib}\right) = 2i \tan^{-1}\left(\frac{b}{a}\right)$ & [8]

$$\cos\left[i \log\left(\frac{a+ib}{a-ib}\right)\right] = \frac{a^2 - b^2}{a^2 + b^2}$$

4. a) If $u = \sin^{-1}\left(\frac{x+y}{\sqrt{x} + \sqrt{y}}\right)$, Prove that [6]

$$x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = \frac{-\sin u \cos 2u}{4 \cos^3 u}$$

b) Using encoding matrix $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$; encode & decode the message [6]

'ALL IS WELL'.

c) Solve the following equations by Gauss Seidal method : [8]

$$10x_1 + x_2 + x_3 = 12$$

$$2x_1 + 10x_2 + x_3 = 13$$

$$2x_1 + 2x_2 + 10x_3 = 14$$

5. a) If $u = e^{xyz} f\left(\frac{xy}{z}\right)$ where, $f\left(\frac{xy}{z}\right)$ is an arbitrary function of $\frac{xy}{z}$, [6]

$$\text{Prove that } x \frac{\partial u}{\partial x} + z \frac{\partial u}{\partial z} = y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 2xyz \cdot u.$$

TURN OVER

- b) Prove that $\sin^5 \theta = \frac{1}{16} (\sin 5\theta - 5\sin 3\theta + 10\sin \theta)$. [6]
- c) i) Prove that $\log(\sec x) = \frac{1}{2} x^2 + \frac{1}{12} x^4 + \dots$ [4]
- ii) Expand $(2x^3 + 7x^2 + x - 1)$ in powers of $(x - 2)$. [4]
6. a) Prove that $\sin^{-1}(\operatorname{cosec} \theta) = \frac{\pi}{2} + i \log \left(\cot \frac{\theta}{2} \right)$. [6]
- b) Find non-singular matrices P & Q such that $A = \begin{bmatrix} 1 & 2 & 3 & 2 \\ 2 & 3 & 5 & 1 \\ 1 & 3 & 4 & 5 \end{bmatrix}$ is [6]
reduced to normal form. Also find its rank.
- c) Obtain the root of $x^3 - x - 1 = 0$ by Regula Falsi Method (Take three [8]
iterations).

TURN OVER