



Applied Physics-I

QP Code:803504

(2 Hours)

[Total Marks : 60

- N.B. :** (1) Question. 1 is compulsory.
 (2) Attempt any **three** questions from the remaining questions No. 2 to 6.
 (3) **Assume** suitable **data** wherever required.
 (4) **Figures** to the **right** indicate **marks**.

1. Attempt any **five** questions from the following-

15

- What are crystal imperfections? Mention any two significance of it.
- Write schrodinger's time dependent and time independent wave equations of matter waves in one dimension and state physical significance of these equations.
- Draw the I-V characteristics of a photo-diode. What is meant by dark current?
- Define super conductivity and critical temperature. Plot the variation of resistance versus temperature in case of superconducting state of the material.
- What is reverberation time? Discuss sabine Formula
- State 'magnetostriction effect.' Mention any two applications of ultrasonic waves.
- Calculate conductivity of a germanium sample if a donar impurity atoms are added to the extent to one part in 10^6 germanium atoms at room temperature.

Assume that only one electron of each atom takes part in conduction process.

Given: Avogadro's number- 6.023×10^{23} atoms/gm - mole

Atomic weight of Ge=72.6

Mobility of electrons = $3800 \text{ cm}^2/\text{volts sec.}$

Density of Ge = 5.32 gm/cm^3

- Describe with necessary theory the Davisson and Germer experiment establishing wave nature of electrons. calculate the de-broglie wavelength of an alpha particle accelerating through a potential difference of 200 volts
 Given- Mass of alpha particle = $6.68 \times 10^{-27} \text{ kg.}$
 - Define the terms drift current and mobility of a charge carriers. Calculate the current product in a germanium sample of area of cross section 1 cm^2 and thickness of 0.01 m , when a potential difference of 2 V is

[Turn Over

applied cross it. Given- The concentration of free electrons in germanium is $2 \times 10^{19}/\text{m}^3$ and mobilities of electrons and holes are $0.36 \text{ m}^2/\text{volts sec}$ and $0.17 \text{ m}^2/\text{volts sec}$ respectively.

3. (a) Draw and explain the unit cell of sodium chloride (NaCl) crystal. Determine effective number of NaCl molecules per unit cell and co-ordination number. 8
- (b) State applications of Hall effect. In a Hall effect experiment a potential difference of $4.5 \mu\text{V}$ is developed across a foil of zinc of thickness 0.02 mm , when a current of 1.5 A is carrying in a direction perpendicular to applied magnetic field of 2 tesla . Calculate 7
- (a) Hall coefficient for zinc
- (b) Concentration of electrons
4. (a) Discuss formation of cooper pairs and energy gap in superconductor on the basis of BCS theory. 5
- (b) State any five factors affecting the acoustics of the building and give the remedies for each. 5
- (c) An ultrasonic pulse of 0.09 MHz sends down towards the sea-bed which returns after 0.55 seconds . The velocity of ultrasonic waves in sea water is 1800 m/sec . Calculate the depth of sea and wavelength of ultrasonic pulse. 5
5. (a) How does the position of Fermi energy level changes with increasing doping concentration in p-type semi-conductors? sketch the diagram. 5
- (b) Explain analysis of crystal structure using Bragg's X ray spectrometer. 5
- (c) Find the minimum energy of neutron confined to a nucleus of size of the order of 10^{-14} m . 5
- Given mass of neutron = $1.675 \times 10^{-27} \text{ kg}$.
6. (a) Calculate the critical radius ratio of an ionic crystal in ligancy -6. What is the maximum size of cation in ligancy-6 configuration, when size of anion is 2.02 \AA ? 5
- (b) What do you mean by group and phase velocity? Show that the de-Broglie group velocity associated with the wave packet is equal to the velocity of the particle. 5
- (c) Explain the formation of potential barrier across the unbiased p-n junction region. 5